2 Prevalence of antibiotic-resistant, toxic metal-tolerant and biofilm- 3 forming bacteria in hospital surroundings

4 Soumitra Nath^{1,2,3,*} , Ahana Sinha¹, Y. Suchitra Singha¹, Ankita Dey¹, Nilakshi Bhattacharjee¹ and Bibhas Deb^{1,2,3}

5
6 ¹Department of Biotechnology, Gurucharan College, Silchar, Assam, India

7 ²Bioinformatics Centre, Gurucharan College, Silchar, Assam, India

8 ³Institutional Biotech Hub, Gurucharan College, Silchar, Assam, India

9 *Correspondence: nath.soumitra1@gmail.com

10
11 Received: April 19, 2020 Accepted: August 31, 2020

Abstract

The emergence and rapid spread of antibiotic-resistant bacteria due to unethical and non-scientific disposal of hospital wastes and clinical by-products caused an alarming environmental concern and associated public health risks. The present study aims to assess the co-selection of antibiotic resistance and heavy metal tolerance by bacteria isolated from hospital effluents. These isolates were also tested for hemolytic activity, pH-tolerance, thermal inactivation, auto-aggregation, cell-surface hydrophobicity and interaction with other bacteria. The study reports the prevalence of antibiotic-resistant and heavy metal tolerant bacteria in clinical effluents and water samples. Most of these isolates were resistant to vancomycin, clindamycin, ampicillin, rifampicin, penicillin-G, methicillin and cefdinir, and evidenced the production of extended-spectrum β -lactamase enzyme. Toxic metals such as cadmium, copper, iron, lead and zinc also exert a selection pressure towards antibiotic resistance. *Pseudomonas aeruginosa* strain GCC_19W3, *Bacillus sp.* strain GCC_19S2 and *Achromobacter spanius* strain GCC_SB1 showed β -hemolysis, evidenced by the complete breakdown of the red blood cells. Highest auto-aggregation was exhibited by *Bacillus sp.* strain GCC_19S2; whereas, maximum cell-surface hydrophobicity was displayed by *P. aeruginosa* strain GCC_19W1. Antagonistic activity by *Stenotrophomonas maltophilia* strain GCC_19W2, *P. aeruginosa* strain GCC_19W3 and strains of *Achromobacter* restricts the growth of other microorganisms by producing some bactericidal substances. The study emphasises undertaking safety measures for the disposal of clinical effluents directly into the environment. The study suggests adopting necessary measures and regulations to restrict the spread of emerging pathogens within the hospital biome and community, which if unnoticed, might pose a significant clinical challenge.

12
13 **Keywords:** Antibiotic-resistant; heavy metals; β -lactamase; β -hemolysis; auto-aggregation

16 Introduction

17 In the environment, antibiotic-resistance emerges in water, soil and sediments through contamination with
18 human and animal wastes that contain antibiotic-resistant bacteria [1]. The discharge of inadequately treated or untreated
19 wastewater from hospitals significantly contributes to the threat of antibiotic resistance [2]. Hospitals use a variety of
20 strategies to protect the patients, but little has been done to safeguard the environment from the contamination due to
21 clinical effluents. The clinical effluents include the wastewater and solid wastes generated from all the medical and non-
22 medical activities. Hospital wastes refer to the infectious, non-hazardous and hazardous liquid waste with sufficient free
23 liquid arising from dental, medical, nursing, pharmaceutical or similar practices [3]. These wastes can be hazardous, toxic
24 and even lethal, because of their high potential for transmitting diseases [4]. Hospital bacterial ecological niches intersect
25 with those in the environment, resulting in public health risks.

26 *Pseudomonas aeruginosa* and *Bacillus sp.* are ubiquitous in hospital wastewater discharges. The infections caused
27 by *P. aeruginosa* become complicated, as they acquire the ability to resist many classes of antibiotics. Some strains of *P.*
28 *aeruginosa* has been reported to survive for months on dry surfaces, and it can persist and grow in contaminated
29 antimicrobial handwash containing triclosan, making it a critical concern for hospital staff [5]. Infections with *Bacillus sp.*
30 are rare, but they are able to produce spores and grow haemolytically on blood agar plates (β -hemolysis). Gram-positive
31 Bacilli can cause many types of infections and are spread to humans in a variety of ways [6,7]. Some of the frequently
32 encountered species include *B. subtilis*, *B. cereus*, *B. licheniformis*, *B. anthracis*, *B. megaterium*, and *B. pumilus*. *Bacillus sp.* may
33 cause ocular infections including conjunctivitis, endophthalmitis, keratitis, dacryocystitis, panophthalmitis and iridocyclitis

[8]. In 2019, a rare case of bloodstream infection with *Bacillus mycoides* in human have been reported in hospital settings of the Netherlands [6]. *Stenotrophomonas maltophilia* has emerged as an important nosocomial pathogen associated with a variety of infections, such as infection of the bloodstream, respiratory tract, urinary tract, bone and joint, endocarditis and meningitis [9]. It is one of the leading opportunistic multidrug-resistance pathogens in hospitals worldwide. The genus *Achromobacter* is an emerging nosocomial pathogen, which the most frequently found in contaminated solutions of dialysis water, demineralized water, water from nebulizers, humidifiers, incubators, extracorporeal circulation systems, poorly preserved heparin flasks and even antiseptic and disinfectant solutions [10,11]. The gram-negative *Cedecea davisae* bacterium has been implicated in causing infection in skin, soft tissue and lung, and in rare cases, they are associated with catheter-related bloodstream infections [12]. Though infections with *C. davisae* or the genus *Cedecea* are rare, treating them is a challenge due to their broad-spectrum antibiotic resistance [13]. The synthesis of extended-spectrum- β -lactamase (ESBL) enzyme is one of the mechanisms by which different bacteria develop antibiotic resistance. These enzymes hydrolyse the β -lactam ring structure of the antibiotic, thereby rendering it inactive [14]. The most common bacteria found to possess ESBL genes include *Escherichia coli* [15], *Klebsiella pneumoniae* [16], *P. aeruginosa* [17], *Bacillus sp.* [18], *Shigella sp.* and *Salmonella sp.*, [19]. Hospital effluents containing various chemical agents, such as antibiotics, toxic heavy metals etc., interact with diverse microbial agents including pathogenic bacteria and lead to an acquisition of selective pressure [20]. The relationship between microbial acquisition of antibiotic resistance and metal tolerance has been extensively studied by many researchers, and both of these resistance genes are found to be located on the same mobile genetic elements [21,22].

It is evident that the resistant bacterial strains gain entrance to the environment via the indiscriminate discharge of untreated or partially treated wastewater from domestic, clinical and industrial sources [14]. To formulate effective practices and policies for combating antibiotic-resistance and to prevent the direct exposure of pathogenic microbes into the environment, it is essential to investigate the biofilm composition, their distribution, degree of antibiotic resistance, and their growth kinetics with change in pH and temperature. Bacteria generally protect itself from external stress such as nutrient starvation or oxidative stress by auto-aggregation mechanism [23]. This property also plays an important role in protection from the host immune system [24]. The molecular mechanisms of bacterial auto-aggregation in aqueous solution may depend on hydrophobic surface properties [25]. The hydrophobic properties favor adhesion to biotic and abiotic surfaces and penetration of host tissues [26]. The efficiency of adhesion, aggregation and biofilm formation by bacteria discharged from hospitals must be evaluated to prevent or reduce the risk of disease transmission and serious damage to the environment.

The bacteria present in hospital effluents regularly interact with the environment, either by synergism or by antagonism relationships [27]. Microorganisms produce antimicrobial substances such as bacteriocin, antibiotic or other toxic components that can directly restrict the growth of other microorganisms. Hussein et al. [28] isolated antibiotic-producing Actinobacteria from the sediment of Ma'in thermal springs (Jordan), which showed an apparent inhibitory activity against the strains of *E. coli*, *P. aeruginosa*, *B. cereus* and *S. aureus*. Many classes of antimicrobial substances have been identified and reported in the literature [29-31].

The present study aims to assess the co-selection of antibiotic resistance and heavy metal tolerance by bacteria isolated from hospital effluents. The ability of these isolates to produce ESBL was evaluated, and their hemolytic activity was also studied. These isolates were also tested for pH tolerance, thermal inactivation, auto-aggregation, cell-surface hydrophobicity and their interaction with other bacteria.

Materials and Methods

Isolation and Characterization of Isolates

Collection of samples

Clinical effluents were collected from four different hospital premises in Silchar, which is situated in Cachar district, in the state of Assam, India (Figure. 1). The sampling sites were drainage area, sewer manhole and clinical garbage dumping site, from where 17 samples were collected in sterile containers and immediately brought to the laboratory for further studies. Drinking water samples were also collected from two hospitals and three nearby tea stalls, to assess the transmission of pathogenic bacteria.

Isolation and characterization of bacteria

Isolation and quantitative computation of bacteria from effluents samples were performed by serial dilution techniques [32]. 100 μ l of samples were spread onto nutrient agar plates and incubated at 37 °C for 24 hours. The total bacterial count was determined by counting the colonies in microprocessor colony counter. Individual distinct colonies were sub-cultured to get pure isolates. Isolates were further identified by colony morphology (shape, structure, colour, pattern and size), Gram's staining and biochemical tests. Biochemical tests include indole production test, methyl red test, Voges-Proskauer test, citrate utilisation test, starch hydrolysis test, catalase activity, oxidase activity and Triple- sugar iron test [33].

Figure 1. Figure caption. If there are multiple panels, they should be listed as: (a) Description of what is contained in the first panel; (b) Description of what is contained in the second panel. Figures should be placed in the main text near to the first time they are cited. A caption on a single line should be centered.

Molecular identification

Genomic DNA was extracted from isolated bacterial cultures [34] and PCR amplification of 16S rDNA gene was achieved by 704F (5'-GTAGCGGTGAAATGCGTAGA-3') and 907R (5'-CCGTC AATTCMTTTRAGTTT-3') [35]. Sequencing of 16S rDNA was carried out at Xcelris Labs Limited, Gujarat, India using ABI 3730xl 96 capillary system using Big Dye Terminator v3.1 kit. The consensus sequence was generated from forward and reverse sequence data, which was used to perform BLAST search. Based on the maximum identity score, first ten sequences were selected, aligned using Clustal-W and a phylogenetic tree was constructed using FastTree [36]. FastTree algorithm improves the topology and the branch lengths of the phylogenetic tree, with maximum-likelihood rearrangements. Geneious R8 software package (Biomatters Ltd., Auckland, New Zealand) was used to perform all the molecular analysis.

Tests concerning the resistance of the strains towards environmental conditions:

Determination of optimum pH

Nutrient agar plates were prepared with varying pH, ranging from pH 2 to pH 8. The test isolates were streaked onto the plates and incubated at 37 °C for 24 hours. Their growth pattern at extreme acidic and basic environment was recorded, and optimum pH was noted.

Determination of thermal death time

Thermal death time at 60 °C and 100 °C was determined by incubating the broth cultures in a temperature-controlled water bath for 1 hour [37]. After every 10 mins, bacterial samples were streaked on freshly prepared nutrient agar plates and incubated for 24 hours at 37 °C.

Test for heavy metal tolerance

Bacterial tolerance to heavy metals (Cadmium (Cd), Copper (Cu), Iron (Fe), Plumbum (Pb) and Zinc (Zn)) was determined by agar dilution method, which was added in the form of cadmium sulfate, copper (II) sulfate, ferrous sulfate, lead acetate and zinc powder [38-40]. The initial concentration of these metal salts in nutrient plates was 100 µg/mL, and bacterial growth was observed by streaking on respective plates. Metal concentration was progressively increased by 50–100 µg/mL on a fresh agar plate, and the minimum inhibitory concentration (MIC) was noted when the isolates failed to grow on respective plates [41]. The experiment was conducted separately for cadmium (Cd), copper (Cu), iron (Fe), Lead (Pb) and Zinc (Zn), taking five replicates at each concentration.

Test concerning putative pathogenicity traits:

Hemolysis test

Freshly cultured bacteria were streaked on blood agar plates containing 5% (v/v) sheep blood and incubated at 30 °C for 24 hours. The blood agar plate was examined for signs of β-hemolysis (clear zone around the colonies), α-hemolysis (green-hued zones around colonies) or γ-hemolysis (no zones around the colonies) [42].

Auto-aggregation assay

Auto-aggregation assays were performed by following the protocol of Basson et al. [43] with minor modifications. The overnight grown bacterial broth was centrifuged at 10,000 rpm for 5 mins at 4 °C, and pellets were collected. Pellets were washed twice by PBS buffer (pH 7.2), re-suspended in 6 mL of PBS buffer, and the initial absorbance was noted at 600 nm. To conduct the auto-aggregation assay, the cell suspension was kept undisturbed for 2 hours at room temperature. After the incubation period, 1 mL from the upper suspension was taken, and the final absorbance was measured at 600 nm. Auto-aggregation was determined by measuring absorbance at 0 hour (initial) and 2 hours (final). The auto-aggregation percentage was measured by the formula: $[(OD_{initial} - OD_{final}) / OD_{initial}] \times 100$ (1)

Cell surface hydrophobicity assay

The 20 hours old bacterial cultures were centrifuged at 12000 rpm for 5 min at 4 °C, and bacterial cell pellets were collected. Cell pellets were washed twice with PBS, and the initial optical density was recorded at 600 nm. The bacterial suspension was then mixed with 1 mL of hydrocarbons (n- Hexadecane and Toluene), vortexed for 2 mins and left undisturbed for 1 hour for phase separation. The aqueous phase was then removed carefully with a micropipette, and the final absorbance was measured at 600 nm. The percentage of cell surface hydrophobicity was determined by the formula: equation 1 [44].

Determination of antibiotic susceptibility and resistance pattern

The isolates were tested against 24 antibiotics using the Kirby-Bauer disc diffusion technique [45]. Antibiotic discs were placed on freshly prepared lawns of each isolate on Mueller-Hinton agar (MHA) plates and incubated at 37 °C for 24–48 hours. The diameter of the inhibition zones was measured, and the strains were classified following the standard antibiotic disc chart. Standard antibiotic discs were procured from ‘HiMedia’ which includes gentamicin (120 µg), vancomycin (30 µg), tetracycline (30 µg), polymixin-B (300 µg), kanamycin (30 µg), ofloxacin (5 µg), co-trimoxazole (25 µg), meropenem (10 µg), ceftriaxone (30 µg), clindamycin (2 µg), ampicillin (10 µg), norfloxacin (10 µg), rifampicin (5 µg), amikacin (30 µg), penicillin-G 10µg, cefdinir (5 µg), ciprofloxacin (5 µg), azithromycin (15 µg), methicilin (5 µg), and streptomycin (10 µg).

Screening for the production of extended-spectrum β-lactamase (ESBL) enzyme

Combined disk method was performed for the phenotypic detection of ESBL producing isolates. In this test, disks containing cefotaxime (30 µg) and ceftazidime (30 µg) with and without clavulanic acid (10 µg) were used. The test isolates were inoculated on the MHA plate and the antibiotic discs with and without clavulanic acid were placed at a distance of 20 mm from each disk. The plates were incubated at 37 °C for 48 hours, and the resulting inhibition zones were measured. ESBL positive isolates were confirmed when the difference of zone diameters between the β-lactam antibiotic disk and the disk containing the antibiotic with clavulanic acid was greater than 5mm [46].

Test concerning competitiveness

Extraction of bacterial metabolites

For extraction of bacterial metabolites, 5 mL of overnight grown broth culture was mixed with an equal volume of ethyl acetate and shaken well in a rotary shaker at 20 rpm for 10 mins. Isolates were centrifuged, and the upper phase has

been transferred into a new flask to evaporate ethyl acetate. Then, the remaining content was mixed with ethyl acetate: acetone: methanol (1:1:1; v/v/v), which is used to evaluate the antagonistic activity [28].

Antagonistic activity of bacterial metabolites against test pathogens

Three strains of bacteria were used as test pathogens to investigate the antagonistic activity of the *Lactobacillus* spp. They are *Bacillus cereus* strain SN_SA (MH482928), *Acinetobacter johnsonii* strain SB_SK (MH482927), *P. aeruginosa* strain GCC_19W1 (MN066610), obtained from the culture collection of Institutional Biotech Hub, Gurucharan College, Silchar. The effectiveness of bacterial metabolites was studied by the well diffusion method [47]. MHA plates were first seeded with the respective test pathogens, and aliquots of 60 µL of the sterile cell-free supernatant were placed in 6 mm diameter wells on MHA plates. After 24 hours of incubation at 37 °C, the diameters of the zones of growth inhibition were classified as sensitive, moderate sensitive or resistant [48]. DMSO was used as control, and the test was performed in triplicate.

Results

Identification of bacteria

It has been observed that isolates GCC_19W1, GCC_19W2 and GCC_19W3, recovered from hospital wastewater forms white, yellow and green colonies, with a jelly-like surface on NA media. Microscopic observation showed that these isolates were gram-negative and rod-shaped. Biochemical test results demonstrated positive results for citrate test, catalase test and TSI test, whereas negative results for indole production test and VP test. Isolate GCC_19W2 showed negative results for MR test, starch hydrolysis and oxidase test (Table 1). Two isolates recovered from solid wastes (GCC_19S1 and GCC_19S2) showed different morphological and biochemical characteristics. GCC_19S1 showed dark pink colour with a jelly-like appearance in NA plates. The isolate was gram-negative rod-shaped and showed negative results for indole production test, MR test, VP test and TSI test. Isolate GCC_19S2 forms white colonies with mucus type appearance on NA plates. Microscopic observation showed the isolate as gram-positive rod-shaped, and biochemical test results evidenced positive results for VP test, catalase test and oxidase test. Drinking water samples were also collected from off-campus hospital area nearby tea stalls. The isolates recovered from hospital drinking water and nearby tea stalls were designated as GCC_SB1 and GCC_SB2 respectively, which appears to be white to light brown and jelly-like appearance in NA plates. Both of these isolates were gram-negative rod-shaped and showed negative test results for indole production, MR, starch hydrolysis and TSI (Table 1). Molecular identification of the isolated bacteria was done by analyzing the 16S rDNA sequence. The BLAST-N algorithm finds regions of similarity between biological sequences present in the database, which demonstrated that the query sequence has 97–99% identity and 100% query coverage with the 16S rDNA of the bacterium recorded in the GenBank. Phylogenetic tree inferred the degree of relatedness between 16S rDNA sequence of the isolates with other closely related sequences retrieved from the database (Figure. 2). Based on these data, the isolates were identified as *P. aeruginosa* strain GCC_19W1 (GenBank accession number: MN066610), *S. maltophilia* strain GCC_19W2 (GenBank accession number: MN066611), *P. aeruginosa* strain GCC_19W3 (GenBank accession number: MT050053), *C. davisae* strain GCC_19S1 (GenBank accession number: MN066609), *Bacillus* sp. strain GCC_19S2 (GenBank accession number: MN066608), *Achromobacter spanius* strain GCC_SB1 (GenBank accession number: MK000623) and *Achromobacter xylosoxidans* strain GCC_SB2 (GenBank accession number: MK000624).

Optimal pH for bacterial growth

Isolated strains of *P. aeruginosa* and *Bacillus* sp. demonstrated optimal growth at pH 7.0–7.5, and their growth in agar plates was not drastically affected with slight increase or decrease in pH. *Achromobacter* sp. exhibited an exponential growth at pH 7.0–8.0, whereas *S. maltophilia* strain GCC_19W2 and *C. davisae* GCC_19S1 exhibited optimal growth at pH 6.5–7.0.

Thermal death time of the isolates

None of the test isolates was able to tolerate the temperature regime of 100 °C. However, when the temperature was lowered to 60 °C, *A. spanius* strain GCC_SB1 showed significant growth for 70 mins. *Bacillus* sp. strain GCC_19S2 and strains of *P. aeruginosa* also demonstrated significant growth at 60 °C up to 50 mins. *S. maltophilia* strain GCC_19W2 and *C. davisae* strain GCC_19S1 showed moderate growth, demonstrating thermal death time of 20 mins at 60 °C.

Figure 2. Figure caption. If there are multiple panels, they should be listed as: (a) Description of what is contained in the first panel; (b) Description of what is contained in the second panel. Figures should be placed in the main text near to the first time they are cited. A caption on a single line should be centered.

220 Table 1. Morphological and biochemical characteristics of isolated bacteria

Bacterial isolates	GCC_19W1	GCC_19W2	GCC_19W3	GCC_19S1	GCC_19S2	GCC_SB1	GCC_SB2
General Characteristics							
Colour	Yellow-green colonies	White to pale yellow colonies	Yellow-green colonies	Dark pink	White colonies	White to light brown	White to light brown
Surface	Jelly	Jelly	Jelly	Jelly	mucus type	Jelly	Jelly
Gram staining	Gram negative	Gram negative	Gram negative	Gram negative	Gram positive	Gram negative	Gram negative
	Rod	Rod	Rod	Rod	Rod	Rod	Rod
Biochemical Characteristics							
Indole	-	-	-	-	-	-	-
MR	+	-	+	-	-	-	-
VP	-	-	-	-	+	+	+
Citrate	+	+	+	+	-	-	+
Starch	+	-	+	+	-	-	-
Hydrolysis	+	-	+	+	-	-	-
Catalase Test	+	+	+	+	+	+	+
Oxidase	+	-	+	+	+	+	+
TSI	+	+	+	-	-	-	-
Suspect organism	<i>Pseudomonas aeruginosa</i>	<i>Stenotrophomonas maltophilia</i>	<i>Pseudomonas aeruginosa</i>	<i>Cedecea davisae</i>	<i>Bacillus sp.</i>	<i>Achromobacter spanius</i>	<i>Achromobacter xylosoxidans</i>

221 “+” indicates positive result and “-” indicates a negative result, MR: , VP: , TSI

222 Tolerance of isolated bacteria towards heavy metals

223 The co-selection of antibiotic resistance and heavy metal tolerance has been observed by all isolates (Table 2). Isolated
 224 strains of *S. maltophilia*, *C. davisae*, *Bacillus sp.* and *Achromobacter sp.* showed maximum growth in NA media supplemented
 225 with zinc. *S. maltophilia* strain GCC_19W2 showed considerable tolerance towards all tested metals, exhibiting a MIC of 3000
 226 ± 54.77 $\mu\text{g/mL}$ for cadmium, 2060 ± 67.82 $\mu\text{g/mL}$ for copper, 3480 ± 58.31 $\mu\text{g/mL}$ for iron and 3980 ± 66.33 $\mu\text{g/mL}$ for lead in
 227 NA media. Isolates *P. aeruginosa* strain GCC_19W1, *C. davisae* strain GCC_19S1 and *A. spanius* strain GCC_SB1 are able to
 228 withstand 4000 $\mu\text{g/mL}$ of lead in NA amended plates. *A. xylosoxidans* strain GCC_SB2 showed maximum tolerance towards
 229 iron with MIC of 4080 ± 48.99 $\mu\text{g/mL}$ of NA. The present study reports a lower tolerance limit of *P. aeruginosa* strain
 230 GCC_19W1 against all the tested metal salts. Copper amendment NA plates showed the minimum growth kinetics, with
 231 MIC ranging from 1000 ± 44.72 to 2160 ± 44 $\mu\text{g/mL}$ of NA.

233 Hemolytic activity of the isolated strains

234 *P. aeruginosa* strain GCC_19W3, *Bacillus sp.* strain GCC_19S2 and *A. spanius* strain GCC_SB1 showed β -hemolysis
 235 (Table 3), which is evidenced by the complete breakdown of the red blood cells leaving a clear zone. The remaining four
 236 isolates showed γ -hemolysis, and no breakdown of the red blood cells occurred.

237 Auto-aggregation of bacterial isolates

238 Bacteria can exhibit different auto-aggregation abilities ranging from high (50% and above), medium (35–50%),
 239 and low (16–35%) [49]. *Bacillus sp.* strain GCC_19S2 reported the highest auto-aggregation ability (53.65 %), followed by *P.*
 240 *aeruginosa* strain GCC_19W3 and *P. aeruginosa* strain GCC_19W1 with auto-aggregation percentage of 38.13 and 41.35
 241 respectively (Table 4). *S. maltophilia* strain GCC_19W2 and strains of *Achromobacter sp.* showed a significant auto-aggregation
 242 ability to form aggregates. The least capability of auto-aggregation by *C. davisae* strain GCC_19S1 (12.29%) demonstrates
 243 their lower efficiency in adhesion and colonisation in the host epithelium.

245 Cell-surface hydrophobicity of isolates

246 The analysis of cell surface hydrophobicity is an essential parameter for determining the colonisation of
 247 microorganisms in the host cells [50]. Among all the isolated strains, maximum hydrophobic interaction was observed by
 248 the strains of *P. aeruginosa* strain GCC_19W1; having hydrophobicity values of 40.11 and 55.09 for n-hexadecane and toluene
 249 respectively. However, the percent hydrophobic index less than 70% cannot be classified as hydrophobic [51]. Therefore,
 250 from the obtained results, all the strains are classified as weak hydrophobic (Table 5).

Table 2. Minimum inhibitory concentration (MIC) of isolated bacteria against Cadmium (Cd), Copper (Cu), Iron (Fe), Plumbum (Pb) and Zinc (Zn), expressed in $\mu\text{g/mL}$ of nutrient agar media

Test isolates	Cadmium Sulfate	Copper Sulfate	Ferrous Sulfate	Lead Acetate	Zinc powder
<i>Pseudomonas aeruginosa</i> strain GCC_19W1	2560 $\pm$ 50.99	1000 $\pm$ 44.72	2520 $\pm$ 37.42	4080 $\pm$ 37.42	5440 $\pm$ 67.82
<i>Stenotrophomonas maltophilia</i> strain GCC_19W2	3000 $\pm$ 54.77	2060 $\pm$ 67.82	3480 $\pm$ 58.31	3980 $\pm$ 66.33	7920 $\pm$ 96.95
<i>Pseudomonas aeruginosa</i> strain GCC_19W3	1480 $\pm$ 20	2040 $\pm$ 74.83	3460 $\pm$ 50.99	2480 $\pm$ 58.31	5020 $\pm$ 66.33
<i>Cedecea davisae</i> strain GCC_19S1	4040 $\pm$ 67.82	1180 $\pm$ 80	3040 $\pm$ 40	4160 $\pm$ 24.49	7940 $\pm$ 50.99
<i>Bacillus</i> sp. strain GCC_19S2	2520 $\pm$ 20	1040 $\pm$ 40	2420 $\pm$ 58.31	3600 $\pm$ 54.77	8020 $\pm$ 37.42
<i>Achromobacter spanius</i> strain GCC_SB1	2600 $\pm$ 54.77	2160 $\pm$ 44	3560 $\pm$ 74.83	4000 $\pm$ 54.77	8140 $\pm$ 132.66
<i>Achromobacter xylosoxidans</i> strain GCC_SB2	2460 $\pm$ 50.99	2000 $\pm$ 54.77	4080 $\pm$ 48.99	2480 $\pm$ 37.42	7720 $\pm$ 37.42

Values are expressed in $\mu\text{g/mL}$ of NA; results are expressed as the means $\pm$ standard deviations of five independent replicates

Table 3. Hemolytic activity of the bacterial isolates

Isolates	Hemolysis type
<i>Pseudomonas aeruginosa</i> strain GCC_19W1	γ
<i>Stenotrophomonas maltophilia</i> strain GCC_19W2	γ
<i>Pseudomonas aeruginosa</i> strain GCC_19W3	β
<i>Cedecea davisae</i> strain GCC_19S1	γ
<i>Bacillus</i> sp. strain GCC_19S2	β
<i>Achromobacter spanius</i> strain GCC_SB1	β
<i>Achromobacter xylosoxidans</i> strain GCC_SB2	γ

Table 4: Auto-aggregation of bacterial isolates

Isolates	Auto-aggregation (%)
<i>Pseudomonas aeruginosa</i> strain GCC_19W1	38.13
<i>Stenotrophomonas maltophilia</i> strain GCC_19W2	30.37
<i>Pseudomonas aeruginosa</i> strain GCC_19W3	41.35
<i>Cedecea davisae</i> strain GCC_19S1	12.29
<i>Bacillus</i> sp. strain GCC_19S2	53.65
<i>Achromobacter spanius</i> strain GCC_SB1	20.04
<i>Achromobacter xylosoxidans</i> strain GCC_SB2	23.07

Antibiotic resistance pattern

The gram-negative strains of *P. aeruginosa* and *S. maltophilia* showed resistance towards vancomycin, co-trimoxazole, clindamycin, ampicillin, rifampicin, penicillin-G, methicillin and cefdinir (Table 6). A similar trend was observed by *Bacillus* sp. strain GCC_19S2, exhibiting their ubiquitous nature, fast adaptation and acquisition of antibiotic resistance. The gram-negative *C. davisae* has been reported to cause infection in the skin and soft tissues, and in some cases associated with bloodstream infections [12]. The prevalence of antibiotic-resistant *C. davisae* outside the hospital environment is rare, but the study cues their outbreak which may cause serious challenge, if unnoticed. The strains of *Achromobacter* sp., isolated from drinking water, were susceptible to all tested antibiotics.

Production of extended-spectrum β -lactamase (ESBL) enzyme

The production of β -lactamase enzyme by gram-negative bacteria has tremendous therapeutic consequences, and if it remains undetected, it might pose a significant clinical challenge. The study reports the prevalence of ESBL positive bacteria by all the gram-negative bacteria (Table 7). These isolates producing the ESBL also conferred resistance towards a group of antibiotics.

Table 5: Cell surface hydrophobicity of the isolated strains

Sample	Hydrophobicity %	
	Hexadecane	Toluene
<i>Pseudomonas aeruginosa</i> strain GCC_19W1	40.11 %	55.09 %
<i>Stenotrophomonas maltophilia</i> strain GCC_19W2	23.72 %	27.30 %
<i>Pseudomonas aeruginosa</i> strain GCC_19W3	38.47 %	43.38 %
<i>Cedecea davisae</i> strain GCC_19S1	35.01 %	30.48 %
<i>Bacillus</i> sp. strain GCC_19S2	12.74 %	29.45 %
<i>Achromobacter spanius</i> strain GCC_SB1	25.15 %	33.72 %
<i>Achromobacter xylosoxidans</i> strain GCC_SB2	18.39 %	24.07 %

Table 6: Antibiotic sensitivity and resistance pattern of isolated bacteria

Antibiotics		<i>Pseudomonas aeruginosa</i> strain GCC_19W1	<i>Stenotrophomonas maltophilia</i> strain GCC_19W2	<i>Pseudomonas aeruginosa</i> strain GCC_19W3	<i>Cedecea davisae</i> strain GCC_19S1	<i>Bacillus</i> sp. strain GCC_19S2	<i>Achromobacter spanius</i> strain GCC_SB1	<i>Achromobacter xylosoxidans</i> strain GCC_SB2
Name	Code							
Gentamicin	HLG 120	24	19	19	21	21	25	27
Vancomycin	VA 30	NI	NI	NI	NI	NI	17	22
Tetracycline	TE 30	NI	16	14	12	9	22	25
Polymyxin-B	PB 300	19	17	15	19	16	16	17
Kanamycin	K 30	17	7	14	13	15	17	22
Ofloxacin	OF 5	27	10	17	18	21	15	15
Co-Trimoxazole	COT 25	4	NI	NI	NI	16	18	10
Meropenem	MRP 10	27	16	21	30	26	30	29
Ceftriaxone	CTR 30	17	NI	10	19	15	15	13
Clindamycin	CD 2	NI	NI	NI	10	NI	11	10
Ampicillin	AMP 10	NI	NI	NI	12	NI	16	15
Norfloxacin	NX 10	15	17	14	21	22	17	21
Rifampicin	RIF 5	NI	6	NI	NI	NI	15	10
Amikacin	AK 30	21	15	15	16	17	20	21
Penicillin-G	P 10	NI	NI	NI	NI	NI	22	28
Streptomycin	S 10	NI	17	9	19	14	25	12
Ciprofloxacin	CIP 5	17	NI	17	29	29	10	16
Azithromycin	AZM 15	17	12	16	14	10	20	23
Methicillin	MET 5	NI	NI	NI	NI	NI	19	12

Cefdinir	CDR 5	NI	NI	NI	NI	NI	18	12
----------	----------	----	----	----	----	----	----	----

Values are expressed in mm; NI: no inhibition zone; diameter of disc = 6 mm

Table 7: Screening for the production of extended-spectrum- β -lactamase (ESBL) enzyme by the test isolates. The test was performed by combined disk method containing cefotaxime (30 μ g) and ceftazidime (30 μ g) with and without clavulanic acid (10 μ g).

Antibiotics		<i>Pseudomonas aeruginosa</i> strain GCC_19W1	<i>Stenotrophomonas maltophilia</i> strain GCC_19W2	<i>Pseudomonas aeruginosa</i> strain GCC_19W3	<i>Cedecea davisae</i> strain GCC_19S1	<i>Bacillus sp.</i> strain GCC_19S2	<i>Achromobacter spanius</i> strain GCC_SB1	<i>Achromobacter xylosoxidans</i> strain GCC_SB2
Name	Code							
Ceftazidime	CAZ 30	12	10	15	15	16	8	18
Ceftazidime/ Clavulanic acid	CAC 30/10	17	8	8	8	14	16	22
Cefotaxime	CTX 30	16	0	17	17	13	18	19
Cefotaxime/ clavulanic acid	CEC 30/10	22	12	17	17	15	28	28

Values are expressed in mm; diameter of disc = 6 mm

Table 8: Antagonistic activity of isolated strains, against three test pathogens

Isolated bacterial strains	Test pathogens		
	<i>Bacillus cereus</i> strain SN_SA	<i>Acinetobacter johnsonii</i> strain SB_SK	<i>Pseudomonas aeruginosa</i> strain GCC_19W1
<i>Pseudomonas aeruginosa</i> strain GCC_19W1	0	0	0
<i>Stenotrophomonas maltophilia</i> strain GCC_19W2	8.67 $\pm$ 0.33	6.33 $\pm$ 0.33	9 $\pm$ 0.58
<i>Pseudomonas aeruginosa</i> strain GCC_19W3	9.33 $\pm$ 0.88	5.67 $\pm$ 0.33	6
<i>Cedecea davisae</i> strain GCC_19S1	0	0	0
<i>Bacillus sp.</i> strain GCC_19S2	0	1.33 $\pm$ 0.67	0
<i>Achromobacter spanius</i> strain GCC_SB1	3.67 $\pm$ 0.33	1.33 $\pm$ 0.67	8.67 $\pm$ 0.33
<i>Achromobacter xylosoxidans</i> strain GCC_SB2	2.33 $\pm$ 0.67	3.67 $\pm$ 0.33	4.67 $\pm$ 0.33
DMSO (control)	0	0	0

Values are expressed in mm; results are expressed as the means $\pm$ standard deviations of three independent replicates

Antagonistic effect of isolated bacterial metabolites on the growth of other bacteria

Antagonism is a widespread bacterial trait, resulting from the production of diffusible inhibitory or antibiotic substances [52,53]. The human opportunist pathogen *S. maltophilia* strain GCC_19W2 demonstrated inhibitory effect against laboratory strains of *P. aeruginosa*, *B. cereus* and *Acinetobacter johnsonii* with a zone of inhibition of 9 $\pm$ 0.58 mm, 8.67 $\pm$ 0.33 mm and 6.33 $\pm$ 0.33 mm respectively. *P. aeruginosa* strain GCC_19W3 and *Achromobacter sp.*, recovered from wastewater and drinking water also evidenced antagonistic activity against the tested bacterial strains (Table 8). *Bacillus sp.* was found to be effective against *A. johnsonii* strain SB_SK, whereas, *C. davisae* strain GCC_19S1 fails to show any inhibition zone. Cross-examination by inoculating the sample from the zone of inhibition on nutrient agar plates showed no growth of lab isolates bacteria, thereby demonstrating their bactericidal activity.

DISCUSSION

In recent years, the emergence and spread of antimicrobial-resistant microorganisms have posed an increasing threat to health care. Hospital-acquired infections due to microorganisms that are resistant to multiple antibiotics have increased morbidity, mortality and the overall cost of healthcare [54]. The present study reports multi-drug resistance pattern of bacteria that may adversely affect the environment, community and sometimes leads to treatment failure. *P. aeruginosa* and *Bacillus* sp. reported in the present study are ubiquitous in hospital wastewater discharges, acquiring a wide range of resistance towards antibiotics, heavy metals and other contaminants [55-57]. The occurrence of *C. davisae* in healthcare-associated infections has also been reported by many researchers [13,12].

The growth of these microbes can be greatly affected by the change in temperature and pH of the environment, and this principle has been adopted as a method of disinfection. It has been observed that, at a temperature regime 100 °C, none of these isolates were able to survive. On the other hand, when the temperature was lowered to 60°C, *A. spanius* strain GCC_SB1, *Bacillus* sp. strain GCC_19S2 and *P. aeruginosa* showed significant growth for 50 mins. Bacterial thermal tolerance can be affected by various factors such as growth temperature [58], pH [59], water activity [60] and growth medium [61]. All the isolates showed varied optimal pH requirements ranging from pH 6-8, whereas extreme acidic and basic pH significantly decreases the bacterial growth pattern. The relation between pH and the bacteriostatic effect has been studied by many researchers [62,63]. Upon entering the cell, the organic acids lower the cytoplasmic pH, resulting in osmotic stress and partial collapse of the transmembrane proton gradient [64,65]. The result of the present study provides an insight into planning successful composting strategies for potentially hazardous contaminants and biological agents. Moreover, proper disposal of hospital effluents should be performed, excessive usage of antibiotics and other chemicals should be restricted, and drinking water must be boiled to restrict or kill any harmful bacteria. Hospital waste contains a large amount of heavy metals such as chromium, cadmium, lead, mercury, zinc etc., as well as organic compounds. These contaminants pollute the environment and pose public health risks [66,67]. In some cases, hospital wastes are treated by incineration, which involves the combustion of organic substances that are contained in waste materials. However, they cannot destroy the metallic components of the waste and concentrate heavy metals into the bottom ash [68].

The virulence nature of these isolates was further accessed by the production of hemolysin. Among all, *P. aeruginosa* strain GCC_19W3, *Bacillus* sp. strain GCC_19S2 and *A. spanius* strain GCC_SB1 showed β -hemolysis, evidenced by the complete breakdown of the red blood cells leaving a clear zone. Hemolysin production among gram-negative bacteria is an indicator of the virulence factors [69]. The ability of auto-aggregation to form aggregates also provides a gateway to colonise to the biotic and abiotic surface and successive penetration to host tissue. Both environmental and pathogenic bacteria exhibit the property of auto-aggregation, generally mediated by self-recognising surface structures, such as proteins and exopolysaccharides [70]. The present study reports moderate auto-aggregation by the strains of *P.* and *Bacillus*, might help in adapting to antibiotics, heavy metals and other environmental contaminants. Depending on the bacteria, the auto-aggregative phenotype may be constitutive or induced under certain conditions, such as stress, oxygen availability or a change in temperature [25]. Bacterial pathogenesis involves the initial adhesion of the bacteria to the intestinal epithelium in the ileum, followed by colonisation of the colon and production of toxin. The toxin impairs epithelial barrier function and ion transport, causing diarrhea [71]. The least capability of auto-aggregation (12.29%) by *C. davisae* strain GCC_19S1 demonstrates their lower efficiency in adhesion and colonisation in the host epithelium. Analysis of cell surface hydrophobicity is an also essential parameter for determining the colonisation of microorganisms in the host cells [50]. Among all the isolated strains, maximum hydrophobic interaction was shown by the strains of *P. aeruginosa*. Similar studies were conducted by Fiorina et al. [72], who isolated bacteria from biofilms of water distribution pipes and colonised catheters from hospitalized patients. Cell surface hydrophobicity is a major parameter for controlling the adhesion of bacteria to surgical tools, medical implants, and also within the environment [73-75]. It also plays an important role in fouling, transmission of infections, and contamination of fruits and vegetables [76,77].

The present study reports the co-selection of antibiotic resistance and heavy metal tolerance by all isolates. The physical linkage of antibiotic resistance and metal resistance is encoded on the plasmid that confers resistance to the bacteria even when only one co-selecting agent (either antibiotics or heavy metals) is present [78-80]. Co-selection of antibiotic resistance and heavy metal tolerance by *P. aeruginosa* is well documented in many studies [81,82,41,83]. *Bacillus* sp. and *Stenotrophomonas* sp. responds to a wide range of environmental contaminants, and exerts co-selection of heavy metal tolerance and antibiotic resistance [41,84,85]. Multidrug-resistant *C. davisae* strain GCC_19S1 reported in the present study showed tolerance to a group of toxic metals. A study conducted by Resende et al. [86] demonstrated that clinical isolates of *C. davisae* exhibiting tolerance towards Nickel (Ni), Zinc (Zn), Chromium(Cr), Copper (Cu), Cadmium(Cd) and Mercury (Hg), acquires a selection pressure towards a group of antibiotics. In another study, Istiaq et al. [87] reported that strains of *A. xylosoxidans* possess genes for Arsenic (As), copper (Cu), zinc (Zn), and cadmium(Cd) resistance, which also display a broad range of resistance against antibiotics belonging to β -lactams, aminoglycosides, cephalosporins, monobactams and even macrolides. Genomic investigation of *S. maltophilia* SJTH1 isolated from the hospitalized patient of PGIMER, Chandigarh, India revealed diverse antibiotic resistance genes, efflux pumps, heavy metal resistance, various transcriptional regulators etc. [88]. Reports are also available on multidrug-resistant *C. davisae* from Indian Hospital [89]. *S. maltophilia* and *Achromobacter xylosoxidans* have been reported as an emerging nosocomial, multidrug-resistant, opportunistic pathogen in

hospital settings, which is also evident in the present study [90-92]. The antibiotics of β -lactam and carbapenems group are the common antibiotics which are used to treat the infections caused by Gram-negative group of bacterial pathogens. Surprisingly, all the gram-negative strains reported in the present study were found to be ESBL positive. The outbreak of multidrug-resistant and extended-spectrum β -lactamase-producing *P. aeruginosa*, *C. davise*, *A. baumannii* and *Proteus mirabilis* has been reported from hospital environment of Italy [93], Ethiopia [94], Korea [95], Malaysia [96], Mexico [97] and United States [98] and India [89]. Various healthcare-associated infections primarily by *Achromobacter xylosoxidans* has been reported in hospital settings of China [92], Tanzania [91], French West Indies [99].

Synergism and antagonism among indigenous bacteria is a common phenomenon. The present study reports antagonistic activity by the *S. maltophilia* strain GCC_19W2, *P. aeruginosa* strain GCC_19W3 and *Achromobacter sp.* against the tested bacterial strains. A similar study was conducted by Afrin, Bhuiyan [100] demonstrated the antagonistic activity of *Bacillus amyloliquefaciens* against multidrug-resistant *Serratia rubidaea*. Another study reports growth inhibition of *Micrococcus luteus* and *Staphylococcus aureus* by antibiotic-producing *Bacillus* and *Lysobacter* [101]. The bactericidal action of the isolated strains suppresses the growth of other tested microorganisms, possibly through competition for nutrients or secreting the inhibitory substances [102]. Microorganisms may also indirectly inhibit or reduce the growth of other microorganisms by changing pH, osmotic pressure and surface tension [100]. Detailed knowledge of synergism and antagonism among the pathogenic and non-pathogenic group of indigenous bacterial isolates is needed to prevent the emergence of drug-resistant strains, chronic toxicity, as well as the potential loss of normal microbiota upon interaction.

CONCLUSION

The present study demonstrates the outrage of different pathogenic bacteria isolated from nosocomial environments. All the isolates showed significant growth at pH 6-8 and are able to survive at 45 °C. *P. aeruginosa* strain GCC_19W3, *Bacillus sp.* strain GCC_19S2 and *A. spanius* strain GCC_SB1 exhibited β -hemolysis, which is an indicator of the virulence factor. A moderate potency of biofilm formation and colonisation of bacteria in the host cells was confirmed by auto-aggregation and cell-surface hydrophobicity analysis. These conducive results ascertain their adhesion to biotic and abiotic surfaces and may result in disease transmission. It has been observed that prolonged exposure of isolates to clinical effluent and toxic chemical might lead to the co-selection of antibiotic resistance and multi-metal tolerance. The gram-negative strains were able to produce β -lactamase enzyme which may pose serious public health risks. Antagonistic activity against other bacteria was evidenced by *S. maltophilia* strain GCC_19W2, *P. aeruginosa* strain GCC_19W3, *A. spanius* strain GCC_SB1 and *A. xylosoxidans* strain GCC_SB2. In conclusion, the study suggests urgent awareness and measures control the spread of emerging pathogens within the hospital biome and community. Further studies are also needed to assess the transmission of disease and the mechanism of pathogenesis.

Acknowledgement

The authors gratefully acknowledge DBT, New Delhi for providing financial assistance to setup Institutional Biotech Hub and Bioinformatics Centre in Gurucharan College, Silchar, India.

Conflict of interest

The authors declare that they have no competing interests.

Funding

The study was not directly funded by any agency or organization; however, laboratory facilities were provided by the institution.

Ethical approval

The study was approved by the Institutional ethical research committee.

Informed consent

Not applicable

CRedit author statement

(Example) AA: Conceptualization, Methodology, Software; BB: Data curation, Writing- Original draft preparation. CC: Visualization, Investigation; DD: Supervision, Writing- Reviewing and Editing. Please consult author guideline for the CRediT taxonomy.

ORCID

Soumitra Nath: 0000-0003-3678-2297

Supplementary Material

Add short descriptions of supplementary material. This material is available online at www.eaht.org.

References

- [1] Marathe NP, Pal C, Gaikwad SS, Jonsson V, Kristiansson E, Larsson DJ. Untreated urban waste contaminates Indian river sediments with resistance genes to last resort antibiotics. *Water Res* 2017;124:388-397
- [2] Medina M-j, Legido-Quigley H, Hsu LY. Antimicrobial Resistance in One Health. In: *Global Health Security*. Springer; 2020. 209-229
- [3] Ali M, Wang W, Chaudhry N, Geng Y. Hospital waste management in developing countries: A mini review. *Waste Manag Res* 2017;35(6):581-592. doi:10.1177/0734242x17691344
- [4] Blass AP, da Costa SEG, de Lima EP, Tortato U, Borges LA. Environmental Performance Measurement in Hospitals: A Bibliometric Review of Literature (1987–2017). In: *Universities and Sustainable Communities: Meeting the Goals of the Agenda 2030*. Springer; 2020. 133-145
- [5] Lanini S, D'Arezzo S, Puro V, Martini L, Imperi F, Piselli P, et al. Molecular epidemiology of a *Pseudomonas aeruginosa* hospital outbreak driven by a contaminated disinfectant-soap dispenser. *PloS one* 2011;6(2):e17064.
- [6] Heidt J, Papaloukas N, Timmerman C. A rare bloodstream infection: *Bacillus mycoides*. *Neth J Med* 2019;77(6):227
- [7] Chougale RA, Kumar PA. Aerobic Microbiological Surveillance of Operation Theatre from a Tertiary Care Hospital, Kolhapur, India. *Int J Curr Microbiol App Sci* 2019;8(10):1210-1215
- [8] Callegan MC, Parkunan SM, Randall CB, Coburn PS, Miller FC, LaGrow AL, et al. The role of pili in *Bacillus cereus* intraocular infection. *Exp Eye Res* 2017;159:69-76
- [9] Kaur P, Gautam V, Tewari R. Distribution of class 1 integrons, *sul1* and *sul2* genes among clinical isolates of *Stenotrophomonas maltophilia* from a tertiary care hospital in North India. *Microb Drug Resist* 2015;21(4):380-385
- [10] Amoureux L, Sauge J, Sarret B, Lhoumeau M, Bajard A, Tetu J, et al. Study of 109 *Achromobacter* spp. isolates from 9 French CF centres reveals the circulation of a multiresistant clone of *A. xylosoxidans* belonging to ST 137. *J Cyst Fibros* 2019;18(6):804-807
- [11] Haviari S, Cassier P, Dananché C, Hulin M, Dauwalder O, Rouvière O, Bertrand X, Perraud M, Bénét T, Vanhems P. Outbreak of *Achromobacter xylosoxidans* and *Ochrobactrum anthropi* infections after prostate biopsies, France, 2014. *Emerg Infect Dis* 2016;22(8):1412
- [12] Ismaael TG, Zamora EM, Khasawneh FA. *Cedecea davisae*'s role in a polymicrobial lung infection in a cystic fibrosis patient. *Case reports in infectious diseases* 2012; 2012
- [13] Sowjanya KV, Sarao MS, Cunningham JM. A Rare Case of *Cedecea Davisae* Bacteremia Presenting as Biliary Sepsis. *Cureus* 2019;11(8)
- [14] Tissera S, Lee SM. Isolation of extended spectrum β -lactamase (ESBL) producing bacteria from urban surface waters in Malaysia. *Malays J Med Sci* 2013;20(3):14
- [15] Dasgupta A, Majumdar T, Bhowmik R. identification and molecular characterisation of ESBL producing uropathogenic *Escherichia coli* strains isolated from a tertiary care hospital of Tripura. *J Evol Med Dent Sci* 2018;7(24):2829-2833
- [16] Akbar SA. The Prevalence of TEM-1 gene causing resistance to beta-lactam antibiotics in *Klebsiella pneumoniae* isolates from clinical samples and plasmid curing. *Int J Med Res Health Sci* 2016;5(11):557-561
- [17] Ellappan K, Narasimha HB, Kumar S. Coexistence of multidrug resistance mechanisms and virulence genes in carbapenem-resistant *Pseudomonas aeruginosa* strains from a tertiary care hospital in South India. *J Glob Antimicrob Re* 2018;12:37-43

- [18] Ain NU, Iftikhar A, Bukhari SS, Abrar S, Hussain S, Haider MH, et al. High frequency and molecular epidemiology of metallo- β -lactamase-producing gram-negative bacilli in a tertiary care hospital in Lahore, Pakistan. *Antimicrob Resist* 2018;7(1):128
- [19] Bialvaei AZ, Kafil HS, Asgharzadeh M, Aghazadeh M, Yousefi M. CTX-M extended-spectrum β -lactamase-producing *Klebsiella* spp, *Salmonella* spp, *Shigella* spp and *Escherichia coli* isolates in Iranian hospitals. *Braz J Microbiol* 2016;47(3):706-711
- [20] Mandal S, Das S, Mandal M. Plasmid Mediated Antibiotic and Heavy Metal Co-Resistance in Bacterial Isolates from Mahananda River Water (Malda, India). *J Transl Med* 2016;6(185):2161-1025
- [21] Di Cesare A, Eckert EM, D'Urso S, Bertoni R, Gillan DC, Wattiez R, et al. Co-occurrence of integrase 1, antibiotic and heavy metal resistance genes in municipal wastewater treatment plants. *Water Res* 2016;94:208-214
- [22] Pathak A, Jaswal R, Chauhan A. Genomic Characterization of a Mercury Resistant *Arthrobacter* sp. H-02-3 Reveals the Presence Heavy Metal and Antibiotic Resistance Determinants. *Front Microbiol* 2020;10:3039
- [23] Dertli E, Mayer MJ, Narbad A. Impact of the exopolysaccharide layer on biofilms, adhesion and resistance to stress in *Lactobacillus johnsonii* FI9785. *BMC Microbiol* 2015;15(1):8
- [24] Galdiero F, Carratelli CR, Nuzzo I, Bentivoglio C, Galdiero M. Phagocytosis of bacterial aggregates by granulocytes. *Eur J Epidemiol* 1988;4(4):456-460
- [25] Farrell A, Quilty B. Substrate-dependent autoaggregation of *Pseudomonas putida* CP1 during the degradation of monochlorophenols and phenol. *J Ind Microbiol Biot* 2002;28(6):316-324
- [26] Heilmann C. Adhesion mechanisms of staphylococci. In: *Bacterial adhesion*. Springer; 2011. 105-123
- [27] Ethica SN, Muslim R, Widyawardhana RBI, Firmansyah A, Imam S. Synergism and Antagonism among Indigenous Hydrolytic Bacteria from Biomedical Wastes for the Generation of Bacterial Consortium Used as Bioremediation Agent. *Int J Environ Sci Dev* 2019;10(12)
- [28] Hussein EI, Jacob JH, Shakhathreh MAK, Al-Razaq MAA, Juhmani A-SF, Cornelison CT. Detection of antibiotic-producing Actinobacteria in the sediment and water of Ma'in thermal springs (Jordan). *Germs* 2018;8(4):191
- [29] Pathma J, Rahul G, Kamaraj K, Subashri R, Sakthivel N. Secondary metabolite production by bacterial antagonists. *J Biol Control* 2011;25(3):165-181
- [30] Abdel-Hamid MS, Anis A, Elbawab RH, Mohammed AA, Orabi SH, Fathalla SI. Distinctive antagonistic role of new *Enterococcus faecium* ER-3M strain and its bacteriocin effect against *Staphylococcus aureus* Pneumonia. *Rend Lincei Sci Fis Nat* 2018;29(3):675-690
- [31] Bindiya E, Tina K, Sasidharan RS, Bhat SG. BaCf3: highly thermostable bacteriocin from *Bacillus amyloliquefaciens* BTSS3 antagonistic on food-borne pathogens. *3 Biotech* 2019;9(4):136
- [32] Yamamoto H, Fukuda I. Microorganism Detection Method. 2016;US Patent 20,160,002,697,
- [33] Holt G, Keong N, Sneath P, Staley J. *Bergey's Manual of Determinative Bacteriology*. Williams and Wilkins Press; 1994
- [34] Green MR, Sambrook J. *Molecular cloning: a laboratory manual*, vol 1. Cold Spring Harbor Laboratory Press; 2012
- [35] Madison JD, Berg EA, Abarca JG, Whitfield SM, Gorbatenko O, Pinto A, et al. Characterization of *Batrachochytrium dendrobatidis* inhibiting bacteria from amphibian populations in Costa Rica. *Front Microbiol* 2017;8:290
- [36] Price MN, Dehal PS, Arkin AP. FastTree 2—approximately maximum-likelihood trees for large alignments. *PloS one* 2010;5(3)
- [37] Juneja VK, Cadavez V, Gonzales-Barron U, Mukhopadhyay S, Friedman M. Effect of pomegranate powder on the heat inactivation of *Escherichia coli* O104: H4 in ground chicken. *Food Control* 2016;70:26-34
- [38] Abou-Shanab R, Van Berkum P, Angle J. Heavy metal resistance and genotypic analysis of metal resistance genes in gram-positive and gram-negative bacteria present in Ni-rich serpentine soil and in the rhizosphere of *Alyssum murale*. *Chemosphere* 2007;68(2):360-367

- [39] Mohamed RM, Abo-Amer AE. Isolation and characterization of heavy-metal resistant microbes from roadside soil and phylloplane. *J Basic Microbiol* 2012;52(1):53-65
- [40] Pandey S, Ghosh PK, Ghosh S, De TK, Maiti TK. Role of heavy metal resistant *Ochrobactrum* sp. and *Bacillus* spp. strains in bioremediation of a rice cultivar and their PGPR like activities. *J Microbiol* 2013;51(1):11-17
- [41] Nath S, Paul P, Roy R, Bhattacharjee S, Deb B. Isolation and identification of metal-tolerant and antibiotic-resistant bacteria from soil samples of Cachar district of Assam, India. *SN Appl Sciences* 2019;1(7):727
- [42] Gerhardt P, Murray R, Costilow R, Nester EW, Wood WA, Krieg NR, et al. *Manual of methods for general bacteriology*; 1981
- [43] Basson A, Flemming L, Chenia H. Evaluation of adherence, hydrophobicity, aggregation, and biofilm development of *Flavobacterium johnsoniae*-like isolates. *Microb Ecol* 2008;55(1):1-14
- [44] Doyle RJ, Rosenberg M. Measurement of microbial adhesion to hydrophobic substrata. In: *Methods in enzymology*, vol 253. Elsevier 1995; 542-550
- [45] Bacter R, Kirby M, Sherris J, Turek M. Antibiotic susceptibility testing by standard single disc diffusion method. *Am J Clin Pathol* 1966;45:493-496
- [46] Ghafourian S, bin Sekawi Z, Sadeghifard N, Mohebi R, Neela VK, Maleki A, et al. The prevalence of ESBLs producing *Klebsiella pneumoniae* isolates in some major hospitals, Iran. *Open Microbiol J* 2011;5:91
- [47] Shakhathreh MAK, Jacob JH, Hussein EI, Masadeh MM, Obeidat SM, Abdul-salam FJ, et al. Microbiological analysis, antimicrobial activity, and heavy-metals content of Jordanian Ma'in hot-springs water. *J Infect Public Heal* 2017;10(6):789-793
- [48] Prabhurajeshwar C, Chandrakanth RK. Probiotic potential of *Lactobacilli* with antagonistic activity against pathogenic strains: An in vitro validation for the production of inhibitory substances. *Biomed J* 2017;40(5):270-283
- [49] Montoro BP, Benomar N, Lavilla Lerma L, Castillo Gutiérrez S, Gálvez A, Abriouel H. Fermented Aloreña table olives as a source of potential probiotic *Lactobacillus pentosus* strains. *Front Microbiol* 2016;7:1583
- [50] Dey DK, Khan I, Kang SC. Anti-bacterial susceptibility profiling of *Weissella confusa* DD_A7 against the multidrug-resistant ESBL-positive *E. coli*. *Microb Pathog* 2019;128:119-130
- [51] Nostro A, Cannatelli M, Crisafi G, Musolino A, Procopio F, Alonzo V. Modifications of hydrophobicity, in vitro adherence and cellular aggregation of *Streptococcus mutans* by *Helichrysum italicum* extract. *Lett Appl Microbiol* 2004;38(5):423-427
- [52] Al-Saedi F, Stones DH, Vaz DP, Krachler AM. Displacement of Pathogens by an Engineered Bacterium Is a Multifactorial Process That Depends on Attachment Competition and Interspecific Antagonism. *Infect Immun* 2016;84(6):1704-1711
- [53] Christensen GJ, Scholz CF, Enghild J, Rohde H, Kilian M, Thürmer A, et al. Antagonism between *Staphylococcus epidermidis* and *Propionibacterium acnes* and its genomic basis. *BMC genomics* 2016;17(1):152
- [54] Barrasa-Villar JI, Aibar-Remón C, Prieto-Andrés P, Mareca-Doñate R, Moliner-Lahoz J. Impact on morbidity, mortality, and length of stay of hospital-acquired infections by resistant microorganisms. *Clin Infect Dis* 2017;65(4):644-652
- [55] Camiade M, Bodilis J, Chaftar N, Riah-Anglet W, Gardères J, Buquet S, et al. Antibiotic resistance patterns of *Pseudomonas* spp. isolated from faecal wastes in the environment and contaminated surface water. *FEMS Microbiol Ecol* 2020;96(2):fiae008.
- [56] Ashfaq M, Khan KN, Rasool S, Mustafa G, Saif-Ur-Rehman M, Nazar MF, et al. Occurrence and ecological risk assessment of fluoroquinolone antibiotics in hospital waste of Lahore, Pakistan. *Environ Toxicol Pharmacol* 2016;42:16-22
- [57] Agarwal S, Kakati B, Khanduri S, Gupta S. Emergence of carbapenem resistant non-fermenting gram-negative bacilli isolated in an ICU of a tertiary care hospital. *J Clin Diagn Res* 2017;11(1):DC04

- 535 [58] Martinez S, López M, Bernardo A. Thermal inactivation of *Enterococcus faecium*: effect of growth temperature and
536 physiological state of microbial cells. *Lett Appl Microbiol* 2003;37(6):475-481
- 537 [59] Alvarez-Ordóñez A, Fernández A, López M, Bernardo A. Relationship between membrane fatty acid composition and
538 heat resistance of acid and cold stressed *Salmonella senftenberg* CECT 4384. *Food Microbiol* 2009;26(3):347-353
- 539 [60] Kieboom J, Kusumaningrum HD, Tempelaars MH, Hazeleger WC, Abee T, Beumer RR. Survival, elongation, and
540 elevated tolerance of *Salmonella enterica* serovar Enteritidis at reduced water activity. *J Food Prot* 2006;69(11):2681-
541 2686
- 542 [61] de Matos RE. Heat resistance of *Listeria monocytogenes* in dairy products as affected by the growth medium. *J Appl*
543 *Microbiol* 1998;84(2):234-239
- 544 [62] Slonczewski JL, Fujisawa M, Dopson M, Krulwich TA. Cytoplasmic pH measurement and homeostasis in bacteria and
545 archaea. *Adv Microb Physiol* 2009;55:1-317
- 546 [63] Saparov SM, Antonenko YN, Pohl P. A new model of weak acid permeation through membranes revisited: does
547 Overton still rule? *Biophys J* 2006;90(11):L86-L88
- 548 [64] Carpenter C, Broadbent JR. External concentration of organic acid anions and pH: key independent variables for
549 studying how organic acids inhibit growth of bacteria in mildly acidic foods. *J Food Sci* 2009;74(1):R12-R15
- 550 [65] Trček J, Mira NP, Jarboe LR. Adaptation and tolerance of bacteria against acetic acid. *Appl Microbiol Biotechnol*
551 2015;99(15):6215-6229
- 552 [66] Zhao L, Zhang F-S, Chen M, Liu Z. Typical pollutants in bottom ashes from a typical medical waste incinerator. *J Hazard*
553 *Mater* 2010;173(1-3):181-185
- 554 [67] Mohajer R, Salehi MH, Mohammadi J, Emami MH, Azarm T. The status of lead and cadmium in soils of high prevalent
555 gastrointestinal cancer region of Isfahan. *J Res Med Sci* 2013;18(3):210
- 556 [68] Anamul HM, Rahman J, Tanvir M. Zn and Ni of bottom ash as a potential diffuse pollutant and their application as
557 'Fine Aggregate'. *J Civ Eng Res* 2012;2(6):64-72
- 558 [69] Barrett TJ, Blake PA. Epidemiological usefulness of changes in hemolytic activity of *Vibrio cholerae* biotype El Tor
559 during the seventh pandemic. *J Clin Microbiol* 1981;13(1):126-129
- 560 [70] Trunk T, Khalil HS, Leo JC. Bacterial autoaggregation. *AIMS microbiol* 2018;4(1):140
- 561 [71] Li Z, Elliott E, Payne J, Isaacs J, Gunning P, O'loughlin EV. Shiga Toxin-Producing *Escherichia coli* Can Impair T84 Cell
562 Structure and Function without Inducing Attaching/Effacing Lesions. *Infect Immun* 1999;67(11):5938-5945
- 563 [72] Fiorina J, Weber M, Block J. Occurrence of lectins and hydrophobicity of bacteria obtained from biofilm of hospital
564 catheters and water pipes *J Appl Microbiol* 2000;89(3):494-500
- 565 [73] Yuan Y, Hays MP, Hardwidge PR, Kim J. Surface characteristics influencing bacterial adhesion to polymeric substrates.
566 *RSC Adv* 2017;7(23):14254-14261
- 567 [74] Khelissa SO, Abdallah M, Jama C, Faille C, Chihib NE. Bacterial contamination and biofilm formation on abiotic surfaces
568 and strategies to overcome their persistence. *J Mater Environ Sci* 2017;8:3326-3346
- 569 [75] Calfee DP. Crisis in hospital-acquired, healthcare-associated infections. *Annu Rev Med* 2012;63:359-371
- 570 [76] Oh JK, Yegin Y, Yang F, Zhang M, Li J, Huang S, et al. The influence of surface chemistry on the kinetics and
571 thermodynamics of bacterial adhesion. *Sci Rep* 2018;8 (1):1-13
- 572 [77] Peleg AY, Hooper DC. Hospital-acquired infections due to gram-negative bacteria. *N Engl J Med* 2010;362(19):1804-
573 1813
- 574 [78] Poole K. At the nexus of antibiotics and metals: the impact of Cu and Zn on antibiotic activity and resistance. *Trends*
575 *Microbiol* 2017;25(10):820-832
- 576 [79] Zhao Y, Cocerva T, Cox S, Tardif S, Su J-Q, Zhu Y-G, et al. Evidence for co-selection of antibiotic resistance genes and
577 mobile genetic elements in metal polluted urban soils. *Sci Total Environ* 2019;656:512-520

- 578 [80] Nath S, Deb B, Sharma I. Isolation of toxic metal-tolerant bacteria from soil and examination of their bioaugmentation
579 potentiality by pot studies in cadmium- and lead-contaminated soil. *Int Microbiol* 2018;21(1):35-45.
- 580 [81] Baker-Austin C, Wright MS, Stepanauskas R, McArthur J. Co-selection of antibiotic and metal resistance. *Trends*
581 *Microbiol* 2006;14 (4):176-182
- 582 [82] Chellaiah ER. Cadmium (heavy metals) bioremediation by *Pseudomonas aeruginosa*: a minireview. *Appl Water Sci*
583 2018;8(6):154
- 584 [83] Nath S, Deb B, Sharma I, Pandey P. Role of cadmium and lead tolerant *pseudomonas aeruginosa* in seedling
585 germination of rice (*Oryza sativa* L.). *J Environ Anal Toxicol* 2014;4(221):2161-0525
- 586 [84] Saranya K, Sundaramanickam A, Shekhar S, Swaminathan S. Biosorption of mercury by *Bacillus thuringiensis*
587 (CASKS3) isolated from mangrove sediments of southeast coast India; 2019
- 588 [85] Piotrowska M, Rzezzycka M, Ostrowski R, Popowska M. Diversity of Antibiotic Resistance Among Bacteria Isolated
589 from Sediments and Water of Carp Farms Located in a Polish Nature Reserve. *Pol J Environ Stud* 2017;26(1)
- 590 [86] Resende JA, Silva VL, Fontes CO, Souza-Filho JA, de Oliveira TLR, Coelho CM, et al. Multidrug-resistance and toxic
591 metal tolerance of medically important bacteria isolated from an aquaculture system. *Microbes and*
592 *environments*:ME12049; 2012
- 593 [87] Istiaq A, Shuvo MSR, Rahman KMJ, Siddique MA, Hossain MA, Sultana M. Adaptation of metal and antibiotic resistant
594 traits in novel β -Proteobacterium *Achromobacter xylosoxidans* BHW-15. *PeerJ* 2019;7:e6537
- 595 [88] Xiong W, Yin C, Wang Y, Lin S, Deng Z, Liang R. Characterization of an efficient estrogen-degrading bacterium
596 *Stenotrophomonas maltophilia* SJTH1 in saline-, alkaline-, heavy metal-contained environments or solid soil and
597 identification of four 17 β -estradiol-oxidizing dehydrogenases. *J Hazard Mater* 2020;385:121616
- 598 [89] Khalid S, Ahmad N, Ali SM, Khan AU. Outbreak of Efficiently Transferred Carbapenem-Resistant bla NDM-Producing
599 Gram-Negative Bacilli Isolated from Neonatal Intensive Care Unit of an Indian Hospital. *Microbial Drug Resistance*;
600 2019
- 601 [90] Ochoa-Sánchez LE, Vinuesa P. Evolutionary genetic analysis uncovers multiple species with distinct habitat preferences
602 and antibiotic resistance phenotypes in the *Stenotrophomonas maltophilia* complex. *Front Microbiol* 2017;8:1548
- 603 [91] Moremi N, Claus H, Hingi M, Vogel U, Mshana SE. Multidrug-resistant *Achromobacter animicus* causing wound
604 infection in a street child in Mwanza, Tanzania. *Diagn Micr Infec Dis* 2017;88(1):58-61
- 605 [92] Liu C, Pan F, Guo J, Yan W, Jin Y, Liu C, et al. Hospital acquired pneumonia due to *Achromobacter* spp. in a geriatric
606 ward in China: clinical characteristic, genome variability, biofilm production, antibiotic resistance and integron in
607 isolated strains. *Front Microbiol* 2016;7:621
- 608 [93] Pagani L, Mantengoli E, Migliavacca R, Nucleo E, Pollini S, Spalla M, et al. Multifocal detection of multidrug-resistant
609 *Pseudomonas aeruginosa* producing the PER-1 extended-spectrum β -lactamase in Northern Italy. *J Clin Microbiol*
610 2004;42(6):2523-2529
- 611 [94] Solomon FB, Wadilo F, Tufa EG, Mitiku M. Extended spectrum and metallo beta-lactamase producing airborne
612 *Pseudomonas aeruginosa* and *Acinetobacter baumannii* in restricted settings of a referral hospital: a neglected
613 condition. *Antimicrob Resist In* 2017;6(1):106
- 614 [95] Kim JY, Park YJ, Kim SI, Kang MW, Lee SO, Lee KY. Nosocomial outbreak by *Proteus mirabilis* producing extended-
615 spectrum β -lactamase VEB-1 in a Korean university hospital. *J Antimicrob Chemother* 2004;54(6):1144-1147
- 616 [96] Lim KT, Yeo CC, Yasin RM, Balan G, Thong KL. Characterization of multidrug-resistant and extended-spectrum β -
617 lactamase-producing *Klebsiella pneumoniae* strains from Malaysian hospitals. *J Med Microbiol* 2009;58(11):1463-
618 1469
- 619 [97] Silva J, Gatica R, Aguilar C, Becerra Z, Garza-Ramos U, Velázquez M, et al. Outbreak of infection with extended-
620 spectrum β -lactamase-producing *Klebsiella pneumoniae* in a Mexican Hospital. *J Clin Microbiol* 2001;39(9):3193-
621 3196

- 622 [98] Hoban DJ, Bouchillon SK, Dowzicky MJ. Antimicrobial susceptibility of extended-spectrum β -lactamase producers and
623 multidrug-resistant *Acinetobacter baumannii* throughout the United States and comparative in vitro activity of
624 tigecycline, a new glycycline antimicrobial. *Diagn Microb Infect Dis* 2007;57(4):423-428
- 625 [99] Marion-Sanchez K, Pailla K, Olive C, Le Coutour X, Derancourt C. *Achromobacter* spp. healthcare associated infections
626 in the French West Indies: a longitudinal study from 2006 to 2016. *BMC Infect Dis* 2019;19(1):795
- 627 [100] Afrin S, Bhuiyan MNI. Antagonistic activity of *Bacillus amyloliquefaciens* subsp. *amyloliquefaciens* against multidrug
628 resistant *Serratia rubidaea*. 2019;bioRxiv:818054
- 629 [101] Falkinham JO, Wall TE, Tanner JR, Tawaha K, Alali FQ, Li C, et al. Proliferation of antibiotic-producing bacteria and
630 concomitant antibiotic production as the basis for the antibiotic activity of Jordan's red soils. *Appl Environ Microbiol*
631 2009;75(9):2735-2741
- 632 [102] Biradar V, Ashwini S, Lachuriye P, Fatima S, Syed A. Isolation of Soil Bacteria for Potential Production of Antibiotics
633 and their Inhibitory Effect on Growth of Pathogens. *Int J Curr Microbiol App Sci* 2016;5(8):514-524